

MA219 – Linear Algebra 2023 Autumn Semester

[You are expected to write proofs / arguments with reasoning provided, in solving these questions.]

Homework Set 5 (*due by Thursday, October 5* in TA's office hours, or previously in class)

Throughout this homework (and this course), \mathbb{F} denotes an arbitrary field.

Question 1. Suppose \mathbb{F} -vector spaces V, W, X have ordered bases (v_1, \dots, v_n) , (w_1, \dots, w_m) , and (x_1, \dots, x_p) for positive integers n, m, p respectively. Write down a basis of the vector space of bilinear maps $: V \times W \rightarrow X$, and prove that it is a basis.

Question 2. Suppose an \mathbb{F} -vector space V has an ordered basis (v_1, \dots, v_n) . For $1 \leq i_0 \leq n$, define the *dual functionals* $\varphi_{i_0} : V \rightarrow \mathbb{F}$ via:

$$\varphi_{i_0} \left(\sum_{i=1}^n c_i v_i \right) := c_{i_0}.$$

Assuming that $\varphi_{i_0} \in V^*$, show that these vectors form a basis of V^* . (This is called the *dual basis*.)

Question 3. Recall the direct product and direct sum (or coproduct) of a set $\{V_i : i \in I\}$ of \mathbb{F} -vector spaces, constructed in class (and studied in the preceding homework set). The goal of this exercise is to show that

$$\left(\bigoplus_{i \in I} V_i \right)^* \cong \prod_{i \in I} V_i^*,$$

by going the 'reverse' way:

- (1) Given $\Phi = (\varphi_i)_{i \in I}$, with each $\varphi_i \in V_i^*$, first show that $(\varphi_i)_{i \in I}$ yields a linear map from $\bigoplus_{i \in I} V_i$ to \mathbb{F} . Let us call this map $T(\Phi)$.
- (2) Show that the assignment $T : \Phi \mapsto T(\Phi)$ is a linear map, from $\prod_{i \in I} V_i^*$ to $\left(\bigoplus_{i \in I} V_i \right)^*$.
- (3) Show that T is one-to-one and onto.

Question 4. (*Quotient spaces – you don't need to submit this solution.*) Suppose V is an \mathbb{F} -vector space, and $W \subset V$ a subspace. Define a relation $v \sim v'$ on V if $v - v' \in W$. Now define the quotient space V/W to be the set of distinct equivalence classes $[v]$, under the relation:

$$a[v] + b[v'] := [av + bv'], \quad a, b \in \mathbb{F}, \quad v, v' \in V.$$

- (1) Verify that \sim above is an equivalence relation on V .
- (2) Check that the addition defined above (setting $a = b = 1$) is associative.
- (3) Check that $\mathbf{0}_{V/W} := [w]$ for $w \in W$ is an (or 'the') additive identity in V/W .
- (4) Assuming that V/W is a vector space under the above operations (which it is!), check that the map $\pi : V \rightarrow V/W$ sending $v \in V$ to $[v]$ is a surjective \mathbb{F} -linear map.
- (5) Prove that the quotient space satisfies the following 'universal property':

Given any \mathbb{F} -vector space Z , and a \mathbb{F} -linear map $\varphi : V \rightarrow Z$ which maps W to $\mathbf{0}_Z$, there exists a unique \mathbb{F} -linear map $\bar{\varphi} : V/W \rightarrow Z$ such that $\varphi = \bar{\varphi} \circ \pi$ (with π as in the previous part).